

Lattice modularity and linear independence

Fabio Cuzzolin

Department of Computer Science, University of Padova, Italy

MSC2000: 06C10,06C05

The notion of *abstract linear independence* on lattices (due to Birkhoff) generalizes the important concept of independence of vectors of a linear space. This relation can be expressed in three different equivalent ways for *atoms* of a semimodular lattice bounded below, but are in general distinct when applied to *arbitrary* elements (when they can fail to be independence relations). Let L be a semimodular lattice with initial element, and define $\mathcal{LI}_1, \mathcal{LI}_2, \mathcal{LI}_3$ in the following way:

1. $\{p_1, \dots, p_n\} \subset L$ are \mathcal{LI}_1 -independent if $p_j \not\leq \bigvee_{i \neq j} p_i \ \forall j$ ($\equiv p_j \wedge \bigvee_{i \neq j} p_i \neq p_j \ \forall j$);
2. $\{p_1, \dots, p_n\} \subset L$ are \mathcal{LI}_2 -independent if $p_k \wedge (p_1 \vee \dots \vee p_{k-1}) = 0$, $k = 2, \dots, n$;
3. $\{p_1, \dots, p_n\} \subset L$ are \mathcal{LI}_3 -independent if $h(p_1 \vee \dots \vee p_n) = h(p_1) + \dots + h(p_n)$, $h(\cdot)$ rank.

The theory of generalized probabilities or *belief functions* provides a hint about how these relations are connected, for it turns out that structured collections of finite domains (*frames*) of belief functions have the algebraic structure of locally Birkhoff lattice. By comparing the behavior of $\mathcal{LI}_1, \mathcal{LI}_2$ and \mathcal{LI}_3 in the landmark cases of frame and projective lattices (since projective lattices are modular) we inferred a number of general results, proving that for a lattice bounded below L :

1. if L is modular then $\mathcal{LI}_2 \Rightarrow \mathcal{LI}_1$;
2. if L is Birkhoff then $\mathcal{LI}_3 \Rightarrow \mathcal{LI}_2$;
3. if L is a modular Birkhoff lattice then $\mathcal{LI}_2 \Rightarrow \mathcal{LI}_3$;
4. if a Birkhoff lattice L is not modular then $\mathcal{LI}_2 \not\equiv \mathcal{LI}_3$.

Therefore we proved an equivalent condition for the modularity of a lattice expressed in terms of equivalence of candidate independence conditions: *a Birkhoff lattice bounded below is modular iff $\mathcal{LI}_2 \equiv \mathcal{LI}_3$.*